



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Eating and rumination activity in 10 cows over 10 days

Braun, Ueli ; Zürcher, S ; Hässig, Michael

Abstract: Eating and rumination activities were evaluated in 10 Brown Swiss cows over 10 days, and the coefficients of variation (CV) were calculated for the investigated variables. A pressure sensor integrated into the noseband of a halter recorded jaw movements during chewing, which allowed the recording of eating and rumination times and the number of regurgitated boluses. The mean CVs ranged from 5.9 to 12.7% and were smaller for rumination (chewing cycles per bolus, 5.9%; daily number of cuds, 8.4%; rumination time, 9.1%) than for eating (eating time, 12.0%; chewing cycles related to eating, 12.7%). We concluded that of eating and rumination variables examined, the number of chewing cycles per regurgitated bolus is the most robust with little variation in individual cows.

DOI: <https://doi.org/10.1016/j.rvsc.2015.05.001>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-112452>

Journal Article

Accepted Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Braun, Ueli; Zürcher, S; Hässig, Michael (2015). Eating and rumination activity in 10 cows over 10 days. Research in Veterinary Science, 101:196-198.

DOI: <https://doi.org/10.1016/j.rvsc.2015.05.001>

1 **Short communication: Eating and rumination activity in 10 cows over 10 days**

2

3 U. Braun*, S. Zürcher, M. Hässig

4

5 *Department of Farm Animals, Vetsuisse Faculty, University of Zurich, Winterthurerstrasse 260,*

6 *CH-8057 Zurich, Switzerland*

7

8 * Corresponding author: Tel.: +41-44-6358241; fax: +41-44-6358904

9 *E-mail address:* ubraun@vetclinics.uzh.ch (U. Braun)

ABSTRACT

Eating and rumination activities were evaluated in 10 **Brown Swiss** cows over 10 days, and the coefficients of variation (CV) were calculated for the investigated variables. A pressure sensor integrated into the noseband of a halter recorded jaw movements during chewing, which allowed the recording of eating and rumination times and the number of regurgitated **boluses**. The mean CVs ranged from 5.9 to 12.7 % and were smaller for rumination (chewing cycles per **bolus**, 5.9 %; daily number of cuds, 8.4 %; rumination time, 9.1 %) than for eating (eating time, 12.0 %; chewing cycles related to eating, 12.7 %). **We concluded that of eating and rumination variables examined, the number of chewing cycles per regurgitated bolus is the most robust with little variation in individual cows.**

Keywords: Cattle; Eating; Rumination; Variation; Coefficient of variation

Several methods have been developed for automated recording of eating and **(or)** rumination **variables** in cattle (Luginbuhl et al., 1987; Matsui and Okubo, 1991; Lindgren, 2009; Schirmann et al., 2009). The technique used in our clinic is based on a pressure sensor integrated into the noseband of a halter that records pressure changes caused by jaw movements. In a first study, ten cows were fitted with a recording halter for 24 hours, and the recordings were compared with those obtained by direct visual observation (Braun et al., 2013). **There was complete or almost complete agreement between data obtained via direct observation and pressure sensor technique,** and phases of eating and rumination were easily distinguished in the automated recordings. In further studies, **we generated reference ranges for eating and rumination in 300 cows (Zürcher, 2014) and investigated these same variables in cows in the peripartum period (Braun et al., 2014).** For experimental purposes, it is convenient to record various activities during a 24-hour period, but we have not critically evaluated whether the pattern established during one day accurately reflects the eating behaviour of a cow. The goal of this study was therefore to record

eating and rumination in ten cows continuously for ten days and analyse the variation coefficients of the various variables.

Ten healthy Brown Swiss cows were used. They were 4.4 ± 1.5 years of age, 12.5 ± 5.0 weeks postpartum and had a daily milk yield of 28.4 ± 2.9 kg. They were kept in tie stalls and fed hay ad libitum during the day. Corn silage was fed several times a day and concentrate twice a day according to the level of production.

Eating and rumination activities were recorded in each cow for ten days using a pressure sensor integrated into the nose band of a halter (Fig. 1, 2) (Braun et al., 2013; Zürcher, 2014).

The coefficient of variation (CV), defined as the ratio of the standard deviation to the mean and expressed as a percentage, was calculated for all variables.

The mean daily eating time of all cows was 316 ± 48 minutes (Table 1) and the mean CV was 12.0 % (4.4 to 18.0 %, Table 2). The mean daily number of chewing cycles (defined as a complete course of movement of the mandible during a single masticatory stroke) during eating was $19,951 \pm 2,968$ (13,881 to 24,465) and the mean CV was 12.7 % (4.4 to 16.8 %). The daily eating times and the daily number of chewing cycles during eating did not differ within cows ($P > 0.05$).

The mean daily rumination time of all cows was 368 ± 54 minutes (Table 1) and the mean CV was 9.1 % (3.2 to 16.4 %, Table 2). The mean daily number of regurgitated boluses was 502 ± 43 and the mean CV was 8.4 % (5.2 to 13.2 %). The mean daily number of chewing cycles per bolus was 49 (38 to 60) and the mean CV was 5.9 % (1.5 to 12.9 %).

The mean overall CVs for the five eating and rumination variables calculated from the ten daily recordings ranged from 4.7 to 13.9 % (Table 2) and were ≤ 10.0 % in seven cows.

Notably, the rumination variables had smaller CVs than the eating variables. Although the external conditions did not change noticeably during the ten-day study period; the eating variables were more strongly affected by external factors than rumination variables. Several

health disorders can affect feed intake in cattle but none occurred in this study. Social interactions that could interfere with feeding are minimal in cows kept in tie stalls. Individual animal factors and ration effects must therefore be considered as causes for the variations. One individual was responsible for feeding the cows and endeavoured to keep the rations consistent; however, the feed was not weighed and small variations from one feeding to another cannot be ruled out. Only three cows (nos. 3, 4, 10) had feeding-related CVs of < 10 %, and these cows also had rumination-related CVs of < 10 %. In the remaining seven cows, these CVs ranged from 12.3 to 18.0 % and 11.5 to 16.8 %, respectively.

Rumination is affected by various factors (DeBoever et al., 1990), most notably by the structure of the forage. Some studies reported shorter rumination times for forage with shorter particle size (Yansari et al., 2004; Yang und Beauchemin, 2006; Adin et al., 2009) but this was not confirmed by another study (Suarez-Mena et al., 2012). Forage with larger particle size may lead to increased eating and rumination times, increasing chewing rate, or both (Yang und Beauchemin, 2006). The cows of this study were fed the same ration throughout the study period, and therefore the effect of particle size on rumination variables variables was neglected. Likewise, the effect of hypocalcaemia and acute mastitis, which have a proven negative impact on rumination (Hansen et al., 2004, Fogsgaard et al., 2013), were also neglected because these diseases did not occur in the present study. Stress and anxiety reduce rumination activity (Herskin, 2004; Bristow und Holmes, 2007), but could also be ruled out. The CVs for rumination variables were much smaller than CVs for the feeding variables in the present study, and a CV of > 10 % occurred for the number of cuds in only five cows (nos. 1, 2, 5, 7, 8), for rumination time in three cows (nos. 2, 5, 7) and for the number of chewing cycles per cud in one cow (2). Of all investigated variables, the latter appears the most robust with little variation in individual cows. The number of chewing cycles per bolus is an excellent criterion to differentiate healthy and diseased cows. Cows with left displacement of the abomasum had

significantly fewer chewing cycles per bolus compared with healthy controls (Braun et al., 2015), and within five days of corrective right-flank omentopexy, the number of chewing cycles increased significantly accompanied by improvement in the overall condition of the cows. Cows with right displacement of the abomasum, caecal dilation and hardware disease also had considerably fewer chewing cycles per bolus than healthy cows (Tschoner, 2013).

Interestingly, three cows with relatively constant eating variables (nos. 3, 4, 10) also had relatively constant rumination variables, and three cows with less constant eating variables (nos. 2, 5, 7) also had less constant rumination variables.

From this study it was concluded that of all eating and rumination variables examined, the number of chewing cycles per regurgitated bolus is the most robust with little variation in individual cows.

References

- Adin, G., Solomon, R., Nikbachat, M., Zenou, A., Yosef, E., Brosh, A., Shabtay, A., Mabjeesh, S. J., Halachmi, I., Miron, J., 2009: Effect of feeding cows in early lactation with diets differing in roughage-neutral detergent fiber content on intake behavior, rumination, and milk production. *Journal of Dairy Science* 92, 3364-3373.
- Braun, U., Trösch, L., Nydegger, F., Hässig, M., 2013: Evaluation of eating and rumination behaviour in cows using a noseband pressure sensor. *BMC Veterinary Research* 9: 164.
- Braun, U., Tschoner, T., Hässig, M., 2014: Evaluation of eating and rumination behaviour using a noseband pressure sensor in cows during the peripartum period. *BMC Veterinary Research* 10: 195.
- Braun, U., Trösch, L., Nuss, K., 2015: Evaluation of eating and rumination behaviour using a noseband pressure sensor in cows after right-flank laparotomy for correction of left displaced abomasum. *Schweizer Archiv für Tierheilkunde* 157, in press.

110 Bristow, D. J., Holmes, D. S., 2007: Cortisol levels and anxiety-related behaviors in cattle.
 111 Physiology & Behavior 90, 626-628.

112 Dado, R. G., Allen, M. S., 1994: Variation in and relationships among feeding, chewing, and
 113 drinking variables for lactating dairy cows. Journal of Dairy Science 77, 132-144.

114 DeBoever, J. L., Andries, J. I., DeBrabander, D., Cottyn, B. G., Buysse, F. X., 1990: Chewing
 115 activity of ruminants as a measure of physical structure. A review of factors affecting it.
 116 Animal Feed Science and Technology 27, 281-291.

117 Fogsgaard, K. K., Røntved, C. M., Sørensen, P., Herskin, M. S., 2012: Sickness behaviour in
 118 dairy cows during Escherichia coli mastitis. Journal of Dairy Science 95, 630-638.

119 Hansen, S. S., Nørgaard, P., Pedersen, C., Jørgensen, R. J., Mellau, L. S., Enemark, J. D., 2003:
 120 The effect of subclinical hypocalcaemia induced by Na₂EDTA on the feed intake and
 121 chewing activity of dairy cows. Veterinary Research Communications 27, 193-205.

122 Herskin, M. S., Munksgaard, L., Ladewig, J., 2004: Effects of acute stressors on nociception,
 123 adrenocortical responses and behavior of dairy cows. Physiology & Behavior 83, 411-420.

124 Lindgren, E., 2009: Validation of rumination measurement equipment and the role of rumination
 125 in dairy cow time budgets. Master Thesis, Swedish University of Agricultural Sciences,
 126 Uppsala.

127 Luginbuhl, J. M., Pond, K. R., Russ, J. C., Burns J. C., 1987: A simple electronic device and
 128 computer interface system for monitoring chewing behavior of stall-fed ruminant animals.
 129 Journal of Dairy Science 70, 1307-1312.

130 Matsui, K., Okubo, T., 1991: A method for quantification of jaw movements suitable for use on
 131 free-ranging cattle. Applied Animal Behaviour Science 32, 107-116.

132 Schirmann, K., Von Keyserlingk, M. A. G., Weary, D. M., Veira, D. M., Heuwieser, W., 2009:
 133 Technical note: Validation of a system for monitoring rumination in dairy cows. Journal of
 134 Dairy Science 92, 6052-6055.

135 Suarez-Mena, F. X., Zanton, G. I., Heinrichs, A. J., 2013: Effect of forage particle length on
 136 rumen fermentation, sorting and chewing activity of late-lactation and non-lactating dairy
 137 cows. *Animal* 7, 272-278.

138 Tschoner, T., 2013: Untersuchungen über das Fressen und Wiederkauen bei kranken Kühen und
 139 bei Kühen um den Zeitpunkt der Geburt. DrMedVet Thesis, Vetsuisse Faculty, University
 140 of Zurich.

141 Yang, W. Z., Beauchemin, K. A., 2006: Effects of physically effective fiber on chewing activity
 142 and ruminal pH of dairy cows fed diets based on barley silage. *Journal of Dairy Science* 89,
 143 217-228.

144 Yansari, A. T., Valizadeh, R., Naserian, A., Christensen, D. A., Yu, P., Sharoodi, F. E., 2004:
 145 Effects of alfalfa particle size and specific gravity on chewing activity, digestibility and
 146 performance of Holstein dairy cows. *Journal of Dairy Science* 87, 3912-3924.

147 Zürcher, S., 2014: Untersuchungen über das Fressen und Wiederkauen von Kühen verschiedener
 148 Rassen mit Hilfe eines Drucksensors im Halfter. DrMedVet Thesis, Vetsuisse Faculty,
 149 University of Zurich.

Legend to Figures

Fig. 1. Brown Swiss cow with a recording halter for the investigation of eating and rumination behaviour. The blue noseband contains the pressure sensor and the brown leather pouch the UBS logger.

Fig. 2. The pressure sensor (in an oil-filled tube, to the right) is attached to the USB logger (in transparent plastic casing, to the left).

157 **Table 1**

158 Coefficients of variation of eating and rumination variables of 10 Brown Swiss cows recorded over 10 days.

159

Variable	Mean \pm sd (range)	Variation coefficient (range)
Eating time (min)	316 \pm 48 (219 – 382)	12.0 (4.8 – 18.0)
Chewing cycles during eating	19,951 \pm 2,968 (13'881 – 24,465)	12.7 (4.4 – 16.8)
Rumination time (min)	368 \pm 54 (262 – 467)	9.1 (3.2 – 16.3)
Number of bolus	502 \pm 43 (436 – 559)	8.4 (5.2 – 13.2)
Chewing cycles per bolus	49 \pm 7 (38 – 60)	5.9 (1.5 – 12.8)

160

161 **Table 2**
 162 Coefficients of variation (expressed as percentage) for eating and rumination variables in 10 Brown Swiss cows calculated from 10 daily
 163 recordings.
 164

Variable	Cow									
	1	2	3	4	5	6	7	8	9	10
Eating time	14.4	15.3	7.3	4.8	18.0	12.3	14.5	12.6	12.5	8.6
Chewing cycles related to eating	14.3	15.1	8.6	4.4	16.2	11.5	16.5	14.1	16.8	9.2
Rumination time	9.5	15.3	7.5	6.0	15.7	7.1	16.4	5.1	5.4	3.2
Number of boluses	10.4	10.3	6.3	5.7	13.2	6.5	10.9	10.0	5.2	6.1
Chewing										

cycles										
per bolus	1.5	12.9	5.0	2.4	6.4	7.8	7.5	2.8	5.4	7.1
Mean CV										
(total)	10.0	13.8	6.9	4.7	13.9	9.0	13.2	8.9	9.1	6.8

165